

TURF FERTILIZER + 0.37% Prodiamine

- For selective pre-emergence control of grass and broadleaf weeds in:

 Established Turf Grasses (excluding golf course putting greens) and lawns

 Landscape Ornamentals (including established perennials and wildflower plantings)

Prodiamine, [N³,N³-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine	0.37%
NERT INGREDIENTS:	
Total:	400.000/

GUARANTEED FERTILIZER ANALYSIS

30-0-10

TOTAL NITROGEN (N)	ı
30.0% Urea Nitrogen*	ı
SOLUBLE POTASH (K ₂ O)	ŀ
TOTAL SULFUR (S)	ŀ
1.4% Free Sulfur	ŀ
DERIVED FROM: Polymer-Coated Sulfur-Coated Urea, Muriate of Potash, Urea	İ
CHLORINE (CI) Max7.5%	ĺ
* 15.0% Controlled Released Urea Nitrogen from Polymer-Coated Sulfur-Coated Urea	ļ

THIS PRODUCT IS NOT INTENDED FOR RETAIL SALE

In the state of Florida, it is recommended that the user of this product follow the Green Industries and Golf Course Best Management Practices found at: http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/BMP_Book_Final.pdf and http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/glfbmp07.pdf

CAUTION KEEP OUT OF REACH OF CHILDREN

See the following section, Precautionary Statements, Hazards to Humans and Domestic Animals, for additional information.

PRECAUTIONARY STATEMENTS • HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using the besides. tobacco or using the toilet.

FIRST AID

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do by a poison control center or doctor. Do not give anything to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center, or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center, or doctor, or when going for treatment.

going for treatment.

Emergency Phone Numbers

(800) 424-9300 CHEMTREC (Transportation and spills)

(800) 222-1222 Poison Control Center ENVIRONMENTAL HAZARDS

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it is toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water hodges or drainage systems. to water bodies or drainage systems.

To not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in

adjacent sites. Do not contaminate water when disposing of equipment wash water.

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation This product is a selective pre-emergence herbicide that provides residual control of many grass and broadleaf

- Established turf grasses (excluding golf course putting greens) and lawns
 Landscape ornamentals
- Established perennials and wildflower plantings

This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when it is activated by at least 1/2 inch of rainfall, irrigation, or shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following

Not for use on plants being grown for; (1) Sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on ornamental plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and

Do not graze or feed livestock forage cut from areas treated with this product. Do not apply aerially. Do not apply to golf course putting greens. Do not apply this product through any type of irrigation system. Do not use on sod farms.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED

CONTROL, CROP INJURY, OR ILLEGAL RESIDUES. RESISTANCE MANAGEMENT

RESISTANCE MANAGEMENT

For resistance management, this product is a Group 3 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 3 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

• Rotate the use of this product or other Group 3 herbicides within a growing access.

- Rotate the use of this product or other Group 3 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
 Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
 Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled ridividuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
 If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance management and/or integrated weed-management recommendations for specific crops and weed

EPA REG. NO. 60063-40-82757

EPA EST. NO. 82757-OH-001 (Lot # Begins with MF) EPA EST. NO. 82757-FL-001 (Lot # Begins with S) EPA EST. NO. 82757-MA-002 (Lot # Begins with H)

TEGRATED WEED PEST MANAGEMENT: Integrate this product into an overall weed pest management ategy whenever the use of an herbicide is required. Practices known to reduce weed development (tillage, crop competition) and herbicide use (weed scouting, proper application timing, banding) should be followed wherever possible. Consult local agricultural and weed authorities for additional IPM strategies established for

WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds:					
Barnyardgrass	Henbit	Purslane, common			
Bluegrass, Annual (Poa annua)	Itchgrass	Pusley, Florida			
Carpetweed	Johnsongrass (from seed)	Rescuegrass ³			
Chickweed, common	Junglerice	Shepherd's Purse ²			
Chickweed, mouse ear (from seed)	Knotweed	Signalgrass, Broadleaf			
Crabgrass (large,smooth)	Kochia	Speedwell, Persian			
Crowfootgrass	Lambsquarter, common	Sprangletop			
Cupgrass, Woolly	Lovegrass	Spurge, Prostrate			
Foxtails, Annual	Panicum (Texas, Fall, Browntop)	Witchgrass			
Goosegrass ¹	Pigweed	Woodsorrel, Yellow (from seed)			

In many areas a single application of 0.65 lb. to 1.5 lb. active ingredient per acre (equal to 4.0 to 9.3 lbs./1000 sq. ft. of this product) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 0.65 lb. to 1.0 lb. a.i. per acre (equal to 4.0 to 6.3 lbs/1000 sq. ft. of this product) followed after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Annual Rate Table. Do not exceed the maximum rate for turf grass species listed in the Maximum

²Applications for this weed should be made in late summer, fall, or winter prior to germination.


Suppression only. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

APPLICATION DIRECTIONS

Apply uniformly with suitable, calibrated application equipment.

ESTABLISHED TURF: This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the Maximum Annual Rates section of this label. Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

CRABGRASS SEED GERMINATION DATES



USE RESTRICTIONS:

- prestrictions apply to the use of this product in turf grasses and lawns:

 DO NOT apply to overseeded turf within 60 days after seeding or until after the second mowing,
 whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.
 - DO NOT cut (harvest) treated sod before 120 days after application.
- DO NOT apply to newly set sod until the following year.

 DO NOT apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

 DO NOT allow people or pets to enter the treated area until the spray has dried.

USE PRECAUTIONS:

The following precautions apply to the use of this product in turf grasses and lawns:

- Application of this product may thin emerged annual bluegrass and newly overseeded grasses.

 Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed

RATES OF APPLICATION

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds. Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

MAXIMUM ANNUAL RATES: This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Maximum Application Rate/Calendar Year of turf fertilizer by turf grass species1							
Turf Species: Lbs. Product/1000 sq.ft. Lbs. a.i./A							
Creeping Bentgrass	4.0	0.65					
Creeping Red Fescue	4.7	0.75					
Buffalograss Kentucky Bluegrass Perennial Ryegrass	6.2	1.0					
Bermudagrass ² Bahiagrass Centipedegrass Seashore Paspalum St. Augustinegrass Tall Fescue (including turf-type) Zoysia	9.3	1.5					

Tall Fescue (including turf-type) Zoysia

These are the maximum rates per calendar year by species limitations.

"May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 0.5 lb. a.i./acre (equal to 3.1 lbs. /1000 sq. ft. of this product). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtali, Goosegrass, and Rescuegrass due to reduced product rates used in sprigging situations.

Do not apply more than 1.5 lbs. a.i. per calendar year per acre (equal to 9.3 lbs./1000 sq. ft. of this product).

Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Rates Table.

The maximum act of application of this product must not exceed the lesser of the applicable rate determine from the Maximum Annual Rates table, and the applicable rate determined in the State Nutrient Restrictions table.

STATE NUTRIENT RESTRICTIONS

STATE NOTRIENT RESTRICTIONS				
State	Readily Available Nitrogen*	Total Nitrogen		
State	Lbs/1,000 sq ft	Lbs/1,000 sq ft		
MD, NH, PA, VA	0.7	0.9		
NJ	0.7	1.0		

NJ

Nitrogen present beyond the amount show as Readily Available must be in the form of Slow-Release nitrogen

WHEN TO APPLY AFTER OVERSEEDING TURF: Do not apply to overseeded turf within 60 days after seeding or until after
the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling
secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION: This product will inhibit the germination of turf species if overseeded too soon

offer periodic forms and interest in the following table before the production of turf species in the second interest in the following table before the production of turf species in the second interest to the following table before the production of turf species in the second interest to the following table before the production of turf species in the second interest to the following table before the production of turf species in the second interest to the following table before the production of turf species in the second interest the second

after application. Follow rates and intervals in the following table below for best overseeding/reseeding results.

LBs Product/Acre	LBs Product/1000 sq.ft.	LBs a.i./A	Months Before Overseeding		
LDS Flouuct/Acre	LBS Product/1000 sq.it.	LDS d.I./A	North	Transition	South
135	3.1	0.50	4	4	4
176	4.0	0.65	5	4	4
203	4.7	0.75	6	5	5
216	5.0	0.80	-	6	6
270	6.2	1.00	-	7	7
308	7.1	1.14	-	-	9
351	8.1	1.30	-	-	10
405	9.3	1.50	-	-	12

Distributed by: Turf Care Supply LLC 50 Pearl Road, Suite 200 • Brunswick, OH 44212

906133

Net Weight 50 LBS (22.67 KG)



TURF FERTILIZER + 0.37% Prodiamine

SPREADER SETTINGS: Spreader settings vary by make and model of spreader. It is recommended that individual spreaders are

Calibrat	ed for the specific product that is it	i be applied. A walking speed o	i 3 miles per nour i	s recommended.
			SPREADER SET	TINGS
	Caroadar Madal	Application Dates1	a i Iba /Aara	(Draduat Iba /1000

	SPREADER SETTINGS			
Spreader Model	Application Rates ¹ a.i. lbs./Acre (Product lbs./1000 sq ft)			
	0.5 (3.1)	0.75 (4.7)	1.0 (6.2)	1.5 (9.3)
LESCO 021820	D 1/2	E 1/2	G 1/2	1 1/2
Lely HR	5.25	6	7	8.5
Vicon 203	15	17	21	27

Application Rates for example only - do not exceed the maximum annual application rate of 1.5 lb a. i./Acre (9.3 lb product/1000

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed, or seed by storage or disposal store this product in its original container in a dry, cool, secured area

PESTICIDE DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions. Wastes resulting from the use of this product must be disposed of on-site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling, if available, or dispose of empty bag into application equipment.

dispose of empty bag in a sanitary landfill or by incineration or if allow

LANDSCAPE ORNAMENTALS

Including Established Perennials and Wildflower Plantings

This product may be applied for residual pre-emergence weed control in ornamentals. **Use rates and When to Apply**Apply in fall and/or spring, 0.65 to 1.5 lbs. active ingredient per acre, which is equal to 4.0 to 9.3 lbs. of Apply in fall and/or spring, 0.65 to 1.5 lbs. active ingredient per acre, which is equal to 4.0 to 9.3 lbs. of this product per 1,000 sq. ft. of ornamental plantings; therefore 20 lbs of this product will treat 2,150 to 5,000 sq. ft. Use the higher rates of application for longer control periods. Sequential applications may be made as long as the total amount of product applied does not exceed the maximum annual application rate of 1.5 lbs. a.i. per acre.

Established Landscape Ornamentals
This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate it into the soil.

Newly Transplanted Landscape Ornamentals
This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

Tolerant Ornamental Species
This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when it is uniformly applied to the soil surface; avoid

to this product. Best results will be obtained when it is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat,

dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned stay out of smoke.			drought or frost damage) some cultivars of listed plants may be sensitive to this product.				
Scientific name Abies spp.	Common name (Tested varieties)	Scientific name Cotoneaster microphyllus	Common name (Tested varieties) Rockspray cotoneaster	Scientific name Juglans spp.*	Common name (Tested varieties) Walnut*	Scientific name Pistacia spp.*	Common name (Tested Varieties) Pistachio*
Abelia grandiflora	Abelia; Sherwood	Crataegus spp.	Hawthorne	Justicia brandegeana	Shrimp plant	Pittosporum rhombifolium	Queensland pittosporum
Acer palmatum	Japanese maple	Crocosmia spp.	Lucifer	Lagerstroemia indica	Crape myrtle	Pittosporum tobira	Japanese pittosporum
Acer platanoides	Norway maple	Cupressus sempervirens	Italian cypress	Lantana montevidensis	Weeping lantana	Podocarpus macrophyllus	Japanese yew
Achillea sp.	Yarrow (King Edward)	Delosperma alba	White trailing ice plant	Lavender spp.	Lavender (Munstead)	Prunus laurocerasus	English laurel
Actinidia chinensis*	Kiwi*	Delphinium spp.	Larkspur (Blue elf)	Leontopodium alpinum	Edelweiss	Prunus spp.*	Almond, Apricot, Nectarine, Peach, Plum, and Prune*
Agapanthus africanus	Lily-of-the-Nile; African Lily	Dianthus deltoides	Maiden pinks (Zing)	Ligustrum amurense	Amur privet	Pseudotsuga menziesii**	Douglas-fir**
Akebia quintata	Five-leaf or Chocolate vine	Dianthus gratianopolitanus	Cheddar pinks	Ligustrum japonicum	Japanese privet	Pyracantha coccinea	Firethorn, scarlet
Allium cernuum	Lady's leek or Nodding onion	Dodonea viscosa	Hop bush	Ligustrum lucidum	Glossy privet (Wax-Leaf)	Pyracantha fortuneana	Firethorn
Anemone hybrida	Japanese anemone	Echinacea purpurea	Coneflower, purple (Magnus)	Ligustrum sinense	Chinese privet (Variegata)	Pyracantha koidzumii	Firethorn
Aquilegia spp.	Columbine (Red and gold)	Elaeagnus pungens	Silverberry	Lilium spp.	Lily (Jazz)	Pyrus spp.	Bradford pear
Arctostaphylos densiflora	Vine Hill manzanita	Euonymus fortunei	Wintercreeper	Liriope muscari	Liriope (Big blue, Lilyturf, Silver mound, Evergreen giant)	Quercus rubra	Red oak
Arctotheca calendula	Cape weed	Euonymus japonica	Japanese spindle tree; Evergreen euonymus	Liriope muscari var. variegata	Liriope, variegated	Quercus shumardii	Shumard oak
Artemesia spp.	Wormwood (Silver mound; Castle)	Euonymus kiautschovica	Spreading euonymus	Liriope spicata	Liriope, creeping	Raphiolepis indica	Indian Hawthorne
Aster sp. or Symphiotrichium sp.	Aster (Bonny blue, Purple dome)	Fatsia japonica	Japanese aralia	Lobelia cardinalis	Cardinal flower	Rhododendron (including Azalea)	Rhododendron & Azalea (Coral Bells, Delaware valley white, Flame creeper, George L. Tabor, PJM, Roseum Elegans, Formosa, Hino-crimson, Wakaebisu, Gumpo)
Aster X frikartii		Forsythia intermedia	Border forsythia	Lonicera japonica	Japanese honeysuckle	Rosa banksiae	Lady Bank's Rose
Athyrium felix-femina	Lady fern	Forsythia suspense	Weeping forsythia	Lonicera tatarica	Tatarian honeysuckle	Rosmarinus officinalis*	Rosemary*
Aucuba japonica	Japanese Aucuba	Forsythia viridissima	Greenstem forsythia	Loropetalum chinense	Burgundy	Rudbeckia spp.	Black-eyed Susan
Begonia spp.	Fibrous bigonia (Hardy Grandis)	Gaillardia spp.	Blanket flower (Goblin)	Lythrum spp.	Loosestrife (Modern Pink)	Rumohra adiantiformis	Leatherleaf Fern
Berberis gladwynesis	Barberry	Gardenia jasminoides	Gardenia, Cape-jasmine	Magnolia spp.**	Magnolia**	Santolina virens	Goldstrum Santolina
Berberis julianae	Wintergreen barberry	Gentiana dahurica	Gentian	Maleophora luteola	Ice plant	Saxifraga spp.	Saxifrage, Purple dome
Berberis mentorensis	Mentor barberry	Geranium cinereum	Cranesbill	Malus spp.*	Crabapple*	Sedum album	Stonecrop
Berberis thunbergii	Japanese barberry	Gladiolus spp.**	Gladiolus	Miscanthus sinensis	Yaku jima, Silberfeder	Sedum caulicola	Stonecrop, likadense
Berberis verriculosa	Warty barberry	Gypsophila repens	Baby's breath	Nandina domestica	Heavenly bamboo	Sedum dasyphyllum	Stonecrop
Bergenia cordifolia		Hedera helix	English Ivy	Narcissus spp.**	Narcissus**	Syzygium paniculatum	Japanese boxcherry
Boltonia asteroides Bougainvillea spp.	Snowbank Bouganvillea	Helianthemum spp. Hemerocallis spp.	Sunrose Daylily (Aztec gold, Stella	Nerium oleander Olea europaea*	Oleander Olive*	Taxus cuspidata Taxus media	Japanese yew Yew
Buddleia davidii	Butterfly-bush (Dwarf Blue,	Hibiscus spp. **	d'Oro, Tender love) Rose of Sharon**	Ophiopogon japonicus**	Mondo Grass**	Teucrium spp.	Germander
Buxus microphylla	Royal Red) Japanese boxwood	Hibiscus spp.	Mallow (Disco belle white)	Osmanthus heterophyllus	Osmanthus; False holly	Thalictrum dipterocarpum	Meadow rue
Callistemon viminalis	Weeping bottlebrush	Hibiscus Rosa-sinensis**	Chinese hibiscus**	Osteospermum fruticosum	(Gulf tide) Trailing African Daisy	Thuja occidentalis	American arborvitae
						<u> </u>	
Calluna vulgaris Campanula carpatica	Scotch heather Tussock bellflower (White	Hosta plantaginea H. sieboldiana	Hosta, Plantain lily Hosta (Searsucker)	Oxydendron luteum Persea americana	Sourwood Avocado*	Trachelospermum asiatum Tsuga canadensis	Star Jasmine Canada hemlock
Campaia V tagliahuana	Clips)						
Campsis X tagliabuana	Trumpet creeper; Trumpet	Houttuynia cordata var.		Photinia fraseri	Frasier's photinia; Redtip	Tulipa spp.	Tulip
Campsis X tagilabuana Carpobrotus edulis		Houttuynia cordata var. variegata Hydrangea macrophylla	Bigleaf hydrangea	Photinia fraseri Physostegia virginiana	Frasier's photinia; Redtip False dragonhead (Vivid)	Tulipa spp. Viburnum japonicum	Tulip Japanese viburnum
<u> </u>	Trumpet creeper; Trumpet flower (Madame Gaten)	variegata	Bigleaf hydrangea Chinese holly**		' '		·
Carpobrotus edulis	Trumpet creeper; Trumpet flower (Madame Gaten) Hottentot fig; Ice plant	variegata Hydrangea macrophylla	· · ·	Physostegia virginiana	False dragonhead (Vivid) Spruce species** (Colorado blue, Norway,	Viburnum japonicum	Japanese viburnum
Carpobrotus edulis Cassia artemisoides Ceanothus rigidus Ceratostigma	Trumpet creeper; Trumpet flower (Madame Gaten) Hottentot fig; Ice plant Feathery Cassia	variegata Hydrangea macrophylla Ilex cornuta**	Chinese holly**	Physostegia virginiana Picea spp.**	False dragonhead (Vivid) Spruce species** (Colorado blue, Norway, etc.)	Viburnum japonicum Viburnum odoratissimum	Japanese viburnum Sweet viburnum
Carpobrotus edulis Cassia artemisoides Ceanothus rigidus Ceratostigma plumbaginoides	Trumpet creeper; Trumpet flower (Madame Gaten) Hottentot fig; Ice plant Feathery Cassia	variegata Hydrangea macrophylla Ilex comuta** Ilex crenata	Chinese holly** Japanese holly	Physostegia virginiana Picea spp.** Pieris japonica	False dragonhead (Vivid) Spruce species** (Colorado blue, Norway, etc.) Lily-of-the-valley shrub	Viburnum japonicum Viburnum odoratissimum Viburnum plicatum	Japanese viburnum Sweet viburnum Japanese snowball
Carpobrotus edulis Cassia artemisoides Ceanothus rigidus Ceratostigma plumbaginoides Chamaecyparis pisitera Chrysanthemum	Trumpet creeper; Trumpet flower (Madame Gaten) Hottentot fig; Ice plant Feathery Cassia Wild Lilac	variegata Hydrangea macrophylla Ilex comuta** Ilex crenata Ilex opaca	Chinese holly** Japanese holly American holly	Physostegia virginiana Picea spp.** Pieris japonica Pinus brutia	False dragonhead (Vivid) Spruce species** (Colorado blue, Norway, etc.) Lily-of-the-valley shrub Calabrian pine	Viburnum japonicum Viburnum odoratissimum Viburnum plicatum Viburnum rigidum	Japanese viburnum Sweet viburnum Japanese snowball Canary Island Viburnum
Carpobrotus edulis Cassia artemisoides Ceanothus rigidus Ceratostigma plumbaginoides Chamaecyparis pisifera Chrysanthemum nipponicum	Trumpet creeper; Trumpet flower (Madame Gaten) Hottentot fig; Ice plant Feathery Cassia Wild Lilac	variegata Hydrangea macrophylla Ilex comuta** Ilex crenata Ilex opaca Ilex pernyi	Chinese holly** Japanese holly American holly Holly	Physostegia virginiana Picea spp.** Pieris japonica Pinus brutia Pinus canariensis	False dragonhead (Vivid) Spruce species** (Colorado blue, Norway, etc.) Lily-of-the-valley shrub Calabrian pine Canary island pine	Viburnum japonicum Viburnum odoratissimum Viburnum plicatum Viburnum rigidum Viburnum tinus	Japanese viburnum Sweet viburnum Japanese snowball Canary Island Viburnum Laurustinus
Carpobrotus edulis Cassia artemisoides Ceanothus rigidus Ceratostigma plumbaginoides Chamaecyparis pisifera	Trumpet creeper; Trumpet flower (Madame Gaten) Hottentot fig; Ice plant Feathery Cassia Wild Lilac False Cypress	variegata Hydrangea macrophylla Ilex comuta** Ilex crenata Ilex opaca Ilex pernyi Ilex vomitoria	Chinese holly** Japanese holly American holly Holly	Physostegia virginiana Picea spp.** Pieris japonica Pinus brutia Pinus canariensis Pinus elliottii	False dragonhead (Vivid) Spruce species** (Colorado blue, Norway, etc.) Lily-of-the-valley shrub Calabrian pine Canary island pine Slash pine	Viburnum japonicum Viburnum odoratissimum Viburnum plicatum Viburnum rigidum Viburnum tinus Viburnum trilobium	Japanese viburnum Sweet viburnum Japanese snowball Canary Island Viburnum Laurustinus Cranberry Bush
Carpobrotus edulis Cassia artemisoides Ceanothus rigidus Ceratostigma plumbaginoides Chamaecyparis pisifera Chrysanthemum nipponicum Cleyera japonica	Trumpet creeper; Trumpet flower (Madame Gaten) Hottentot fig; Ice plant Feathery Cassia Wild Lilac False Cypress	variegata Hydrangea macrophylla Ilex comuta** Ilex crenata Ilex opaca Ilex pernyi Ilex vomitoria Inula ensifolia	Chinese holly** Japanese holly American holly Holly Yaupon holly	Physostegia virginiana Picea spp.** Pieris japonica Pinus brutia Pinus canariensis Pinus elliottii Pinus halepensis	False dragonhead (Vivid) Spruce species** (Colorado blue, Norway, etc.) Lily-of-the-valley shrub Calabrian pine Canary island pine Slash pine Aleppo pine	Viburnum japonicum Viburnum odoratissimum Viburnum plicatum Viburnum rigidum Viburnum tinus Viburnum trilobium Viburnum wrightii	Japanese viburnum Sweet viburnum Japanese snowball Canary Island Viburnum Laurustinus Cranberry Bush Leatherleaf Viburnum
Carpobrotus edulis Cassia artemisoides Ceanothus rigidus Ceratostigma plumbaginoides Chamaecyparis pisifera Chrysanthemum nipponicum Cleyera japonica Citrus spp. *	Trumpet creeper; Trumpet flower (Madame Gaten) Hottentot fig; Ice plant Feathery Cassia Wild Lilac False Cypress Cleyera Citrus species* Coreopsis; Calliopsis (Early	variegata Hydrangea macrophylla Ilex cornuta** Ilex crenata Ilex opaca Ilex pernyi Ilex vomitoria Inula ensifolia Iris ensata	Chinese holly** Japanese holly American holly Holly Yaupon holly Sword-leaved iris (Jodlesong)	Physostegia virginiana Picea spp.** Pieris japonica Pinus brutia Pinus canariensis Pinus elliottii Pinus halepensis Pinus nigra	False dragonhead (Vivid) Spruce species** (Colorado blue, Norway, etc.) Lily-of-the-valley shrub Calabrian pine Canary island pine Slash pine Aleppo pine Austrian black pine	Viburnum japonicum Viburnum odoratissimum Viburnum plicatum Viburnum rigidum Viburnum tinus Viburnum trilobium Viburnum wrightii Vinca major	Japanese viburnum Sweet viburnum Japanese snowball Canary Island Viburnum Laurustinus Cranberry Bush Leatherleaf Viburnum Greater periwinkle, Vinca
Carpobrotus edulis Cassia artemisoides Ceanothus rigidus Ceratostigma plumbaginoides Chamaecyparis pisifera Chrysanthemum nipiopnicum Cleyera japonica Citrus spp. * Coreopsis spp. Cornus florida	Trumpet creeper; Trumpet flower (Madame Gaten) Hottentot fig; Ice plant Feathery Cassia Wild Lilac False Cypress Cleyera Citrus species* Coreopsis; Calliopsis (Early sunrise, Moonbeam)	variegata Hydrangea macrophylla Ilex comuta** Ilex crenata Ilex opaca Ilex pernyi Ilex vomitoria Inula ensifolia Iris ensata Iris siberica	Chinese holly** Japanese holly American holly Holly Yaupon holly Sword-leaved iris (Jodlesong) Siberian iris (Cabernet)	Physostegia virginiana Picea spp.** Pieris Japonica Pinus brutia Pinus canariensis Pinus elliotii Pinus halepensis Pinus nigra Pinus palustris	False dragonhead (Vivid) Spruce species** (Colorado blue, Norway, etc.) Lily-of-the-valley shrub Calabrian pine Canary island pine Slash pine Aleppo pine Austrian black pine Longleaf pine	Viburnum japonicum Viburnum odoratissimum Viburnum plicatum Viburnum rigidum Viburnum tinus Viburnum tirilobium Viburnum wrightii Vinca major Vinca minor	Japanese viburnum Sweet viburnum Japanese snowball Canary Island Viburnum Laurustinus Cranberry Bush Leatherleaf Viburnum Greater periwinkle, Vinca Common periwinkle
Carpobrotus edulis Cassia artemisoides Ceanothus rigidus Ceratostigma plumbaginoides Chamaecyparis pisifera Chrysanthemum nipponicum Cleyera japonica Citrus spp. * Coreopsis spp. Cornus florida Cornus stolonifera	Trumpet creeper; Trumpet flower (Madame Gaten) Hottentot fig; Ice plant Feathery Cassia Wild Lilac False Cypress Cleyera Citrus species* Coreopsis; Calliopsis (Early sunrise, Moonbeam) Flowering dogwood	variegata Hydrangea macrophylla Ilex cornuta** Ilex crenata Ilex opaca Ilex pernyi Ilex vomitoria Inula ensifolia Iris ensata Iris siberica Iris spp.	Chinese holly** Japanese holly American holly Holly Yaupon holly Sword-leaved iris (Jodlesong) Siberian iris (Cabernet) Iris ** Winter jasmine Chinese juniper (Nick's	Physostegia virginiana Picea spp.** Pieris japonica Pinus brutia Pinus canariensis Pinus elliottii Pinus halepensis Pinus nigra Pinus palustris Pinus radiate	False dragonhead (Vivid) Spruce species** (Colorado blue, Norway, etc.) Lily-of-the-valley shrub Calabrian pine Canary island pine Slash pine Aleppo pine Austrian black pine Longleaf pine Monterey pine	Viburnum japonicum Viburnum odoratissimum Viburnum plicatum Viburnum rigidum Viburnum tinus Viburnum trilobium Viburnum wrightii Vinca major Vinca minor Vitis spp.*	Japanese viburnum Sweet viburnum Japanese snowball Canary Island Viburnum Laurustinus Cranberry Bush Leatherleaf Viburnum Greater periwinkle, Vinca Common periwinkle Grape*
Carpobrotus edulis Cassia artemisoides Ceanothus rigidus Ceratostigma plumbaginoides Chamaecyparis pisifera Chrysanthemum nipponicum Cleyera japonica Citrus spp. * Coreopsis spp. Comus florida Cornus stolonifera Cortaderia selloana Cotoneaster apiculatus	Trumpet creeper; Trumpet flower (Madame Gaten) Hottentot fig; Ice plant Feathery Cassia Wild Lilac False Cypress Cleyera Citrus species* Coreopsis; Calliopsis (Early sunrise, Moonbeam) Flowering dogwood American dogwood Pampas grass Cranberry cotoneaster	variegata Hydrangea macrophylla Ilex cornuta** Ilex crenata Ilex opaca Ilex pernyi Ilex vomitoria Inula ensifolia Iris ensata Iris siberica Iris spp. Jasminium nudiflorum	Chinese holly** Japanese holly American holly Holly Yaupon holly Sword-leaved iris (Jodlesong) Siberian iris (Cabernet) Iris ** Winter jasmine	Physostegia virginiana Picea spp.** Pieris japonica Pinus brutia Pinus canariensis Pinus elliottii Pinus halepensis Pinus nigra Pinus palustris Pinus radiate Pinus strobus	False dragonhead (Vivid) Spruce species** (Colorado blue, Norway, etc.) Lily-of-the-valley shrub Calabrian pine Canary island pine Slash pine Aleppo pine Austrian black pine Longleaf pine Monterey pine Eastern white pine Scotch pine Loblolly pine	Viburnum japonicum Viburnum odoratissimum Viburnum plicatum Viburnum rigidum Viburnum tinus Viburnum trilobium Viburnum wrightii Vinca major Vinca minor Vitis spp.* Weigela florida Yucca aloifolia	Japanese viburnum Sweet viburnum Japanese snowball Canary Island Viburnum Laurustinus Cranberry Bush Leatherleaf Viburnum Greater periwinkle, Vinca Common periwinkle Grape* Old fashioned Weigela Spanish bayonet Yucca, Adam's needle
Carpobrotus edulis Cassia artemisoides Ceanothus rigidus Ceratostigma plumbaginoides Chamaecyparis pisifera Chrysanthemum nipponicum Cleyera japonica Citrus spp. * Coreopsis spp. Comus florida Cornus stolonifera Cordaderia selloana	Trumpet creeper; Trumpet flower (Madame Gaten) Hottentot fig; Ice plant Feathery Cassia Wild Lilac False Cypress Cleyera Citrus species* Coreopsis; Calliopsis (Early sunrise, Moonbeam) Flowering dogwood American dogwood Pampas grass	variegata Hydrangea macrophylla Ilex crenata Ilex crenata Ilex opaca Ilex pernyi Ilex vomitoria Inula ensifolia Iris ensata Iris siberica Iris spp. Jasminium nudiflorum Juniperus chinensis	Chinese holly** Japanese holly American holly Holly Yaupon holly Sword-leaved iris (Jodlesong) Siberian iris (Cabernet) Iris ** Winter jasmine Chinese juniper (Nick's compact)	Physostegia virginiana Picea spp.** Pieris japonica Pinus brutia Pinus canariensis Pinus elliottii Pinus halepensis Pinus nigra Pinus palustris Pinus radiate Pinus strobus Pinus sylvestris	False dragonhead (Vivid) Spruce species** (Colorado blue, Norway, etc.) Lily-of-the-valley shrub Calabrian pine Canary island pine Slash pine Aleppo pine Austrian black pine Longleaf pine Monterey pine Eastern white pine Scotch pine	Viburnum japonicum Viburnum odoratissimum Viburnum plicatum Viburnum rigidum Viburnum tinus Viburnum trilobium Viburnum wrightii Vinca major Vinca minor Vitis spp.* Weigela florida Yucca aloifolia	Japanese viburnum Sweet viburnum Japanese snowball Canary Island Viburnum Laurustinus Cranberry Bush Leatherleaf Viburnum Greater periwinkle, Vinca Common periwinkle Grape* Old fashioned Weigela Spanish bayonet Yucca, Adam's needle

WARRANTY AND LIMITATION OF DAMAGES

WARRANTY AND LIMITATION OF DAMAGES

Conditions of sale: To the extent consistent with applicable law, Turf Care Supply LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to Turf Care Supply LLC. DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED. To the extent consistent with applicable law, Turf Care Supply LLC. SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, AND TURF CARE SUPPLY LLC'S SOLE LIABILITY AND BUYER AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER AND USER ACKNOWLEDGE AND ASSUME ALL RISKS AND LIABILITY RESULTING FROM HANDLING, STORAGE AND USE OF THIS PRODUCT. TURF CARE SUPPLY LLC DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT.

EPA REG. NO. 60063-40-82757

EPA EST. NO. 82757-OH-001 (Lot # Begins with MF) EPA EST. NO. 82757-FL-001 (Lot # Begins with S) EPA EST. NO. 82757-MA-002 (Lot # Begins with H)

906133 Net Weight 50 LBS (22.67 KG)

Distributed by: Turf Care Supply LLC 50 Pearl Road, Suite 200 Brunswick, OH 44212

